

# FTM (Fluid Thioglycollate Medium) (KP)



Medium used for the cultivation and enumeration of aerobic and anaerobic organisms and for the sterility testing of biologics.

\*Equally use with FTM (Fluid Thioglycollate Medium) Modified (MB-F0777), USP, EP.

## • CONTENTS (Liter)

Yeast Extract	5.0 g
Casein Peptone	15.0 g
Dextrose	5.0 g
Sodium Chloride	2.5 g
L-Cystine	0.5 g
Sodium Thioglycollate	0.5 g
Resazurin	0.001 g
Agar	0.75 g
Final pH = 7.1 ± 0.2 at 25°C	

## • PROCEDURE

Suspend 29.25 G of powder in 1 L of distilled or deionized water. Heat to boiling until completely dissolved. Pour into tubes. Sterilize by autoclave at 121°C for 15 minutes. Cool to 45 - 50°C in water bath. Store fluid thioglycollate media at 2 - 25°C. If more than 30% of the medium is pink prior to use, reheat once (100°C) to drive off absorbed oxygen.

## • INTERPRETATION

FTM (Fluid Thioglycollate Medium) (KP) is a medium used for the cultivation and enumeration of aerobic and anaerobic organisms and for the sterility testing of biologics. Yeast extract and casein peptone provide carbon, nitrogen, vitamins and minerals. Dextrose is a source of carbohydrate. Sodium chloride maintains the osmotic balance. L-Cystine and sodium thioglycollate are the reducing agents. Resazurin is an oxidation-reduction indicator which becomes pink when oxidized and colorless when reduced. Agar is the solidifying agent. Low concentration of agar interrupts the diffusion of oxygen in the medium.

## • TECHNIC

Inoculate the specimen using a sterile needle to the middle of the medium. Incubate at 30 - 35°C for 48 ± 3 hours up to 3 days. Slightly loosen the caps. Refer appropriate references for recommended test procedure.

## • QUALITY CONTROL FOR USE

### Dehydrated medium

Appearance: free-flowing, homogeneous

Color: beige

### Prepared medium

Appearance : clear to slightly opalescent

Color: light amber with upper medium pink

Incubation conditions: 30 - 35°C / 48 ± 3 hours up to 3 days / loosen the caps

Microorganism	ATCC	Inoculum CFU	Growth
<i>Staphylococcus aureus</i>	6538	50-100	good
<i>Bacillus subtilis</i>	6633	50-100	good
<i>Pseudomonas aeruginosa</i>	9027	50-100	good
<i>Clostridium perfringens</i>	13124	50-100	good
<i>Clostridium sporogenes</i>	19404	50-100	good

## • WARNING AND PRECAUTIONS

Check the label before opening the container and follow appropriate microbiological laboratory procedures. Wear protective gloves, protective clothing, safety glasses, and face protection. Safety instructions can be found in Material Safety Data Sheet (MSDS).

## • PERFORMANCE AND EVALUATION

Performance of the medium can be expected when used according to the directions on the label within the recommended temperature.

## • STORAGE AND SHELF LIFE

The powder is very hygroscopic. Store in the provided container according to the temperature indicated on the container surface and store the prepared medium at a temperature of 2 to 8°C. Use within the expiration date indicated on the label and store in a dry and well-ventilated place. The product can be used until the stated expiration date.

## • REFERENCES

1. Quastel and Stephenson. 1926. J. Biochem. 20:1125.
2. Falk, Bucca and Simmons. 1939. J. Bacteriol. 37:121.
3. Brewer. 1940. JAMA 115:598.
4. Marshall, Ginnish and Luxen. 1940. Proc. Soc. Exp. Biol. Med. 43:672.
5. Nungester, Hood and Warren. 1943. Proc. Soc. Exp. Biol. Med. 52:287.
6. Refer to the KP, USP and EP.

## • PACKAGE

Cat. No : MB-F1026P FTM (Fluid Thioglycollate Medium) (KP)	500 G
---	-------